

**BIOMARK Laboratories-INDIA**

[www.biomarklabs.com](http://www.biomarklabs.com)

**TECHNICAL SHEET**

<b>BS011</b>	<b>ANTIBIOTIC ASSAY MEDIUM NO.12(NYSTATIN ASSAY AGAR)</b>					
<b>Formula</b>						
<b>Ingredients:</b>			<b>gms/lit.</b>			
Peptic digest of animal tissue			10.00			
Yeast extract			5.00			
Meat extract B#			2.50			
Dextrose			10.00			
Sodium chloride			10.00			
Agar			25.00			
# Equivalent to Beef extract						
Final pH (at 25°C) : 6.0 ± 0.2						
<b>Directions :</b>						
Suspend 62.5 gms in 1000ml. distilled water. Heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
<b>Principle :</b>						
Nutrients and growth factors are supplied by the ingredients like Peptic digest of animal tissues, Yeast extract and Beef extract etc. Sodium chloride maintains the osmotic equilibrium. Dextrose as a sugar is also supplemented as a carbon and energy source. Agar is the solidifying agent.						
<b>QC Tests - (I) Dehydrated Medium</b>						
Colour :		Cream to light yellow				
Appearance :		Homogeneous Free Flowing powder				
<b>(II) Rehydrated medium</b>						
pH (post autoclaving/heating):		6.0 ± 0.2				
Colour (post autoclaving/heating) :		Cream to yellow				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent				
<b>(III) Q.C. Test Microbiological</b>						
Cultural characteristics observed after 18-24 hrs. at 35-37°C.						
MICROORGANISM (ATCC)		GROWTH	ANTIBIOTICS ASSAYED			
Saccharomyces cerevisiae (2601 )		Luxuriant	Amphotericin B, Nystatin			
<b>Precautions :</b>		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
<b>Use:</b>		It is used for microbiological assay of Amphotericin B and Nystatin using Saccharomyces cerevisiae as per USP/IP.				
<b>Storage:</b>		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
<b>Packing :</b>		500 gm bottle				
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>BS011</b>		62.5 g/l	8.0 L	6.0±0.2	Nil	121°C / 15 minutes

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.