

TECHNICAL SHEET

BW298	PLATE COUNT AGAR					
Formula						
Ingredients :			gms/lit.			
Tryptone			5.00			
Yeast extract			2.50			
Dextrose			1.00			
Sodium chloride			6.50			
Agar			15.00			
Final pH (at 25°C) : 7.0 ± 0.2						
Directions :						
Suspend 30.0 gms in 1000 ml. distilled water. Boil to dissolve the medium completely. Dispense as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
Principle :						
Plate count agar contains Casein enzymic hydrolysate and Yeast extract which provide the carbon and nitrogen sources required for growth of a wide variety of organisms. Dextrose is a source of fermentable carbohydrate (energy source). Agar is a solidifying agent.						
QC Tests – (I) Dehydrated Medium						
Colour :		Cream to light yellow				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		7.0 ± 0.2				
Colour (post autoclaving/heating) :		Cream to light yellow				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 18 – 24 hrs at 35 – 37°C.						
MICROORGANISM (ATCC)		GROWTH				
Bacillus subtilis (6633)		Luxuriant				
Escherichia coli (25922)		Luxuriant				
Lactobacillus casei (9595)		Luxuriant				
Staphylococcus aureus (25923)		Luxuriant				
Enterococcus faecalis (29212)		Luxuriant				
Streptococcus pyogenes (19615)		Luxuriant				
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For determination of plate counts of microorganisms in foods, water and wastewater.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
BW298		30.0g/l	16.66 L	7.0 ± 0.2	NIL	121°C / 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.