

TECHNICAL SHEET

PP001	Bi.G.G.Y. Agar Plate (Nickerson Agar Plate)			
Formula				
Ingredients:	gms/lit			
Yeast extract	1.00			
Glycine	10.00			
Dextrose	10.00			
Bismuth ammonium citrate	5.00			
Sodium sulphite	3.00			
Agar	16.00			
Final pH (at 25°C): 6.8 ± 0.2				
Directions:				
Label the ready to use plate (PP001). Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.				
Principle:				
Bismuth ammonium citrate and Sodium sulphite together act as selective agents for Candida species suppressing bacterial growth, at the same time indicating substrate reduction to yield bismuth sulphite which helps to presumptively identify the culture of Candida species. Yeast extract, dextrose and glycine serve as nutrients. Growth on an acidic or neutral medium containing bismuth sulphite produced black colonies because of the extra cellular reaction of the bismuth sulphite to bismuth sulphide.				
(I) QC Tests				
pH:	6.8 ± 0.2			
Color:	White to Off white coloured medium.			
Appearance:	Sterile Bi.G.G.Y.(Nickerson) Agar 85mm disposable plates.			
(II) Sterility test				
Passes release criteria				
(III) Q.C. Test Microbiological				
Cultural characteristics observed after incubation at 35-37°C for 18-24 hours.				
MICROORGANISM (ATCC)	Inoculum	GROWTH	RECOVERY	COLONY MORPHOLOGY
Candida albicans 10231	50-100	luxuriant	>=50%	smooth, circular intensely brown black, no color diffusion
Candida kruisei 24408	50-100	luxuriant	>=50%	large flat, wrinkled silvery brown, black colonies with brown peripheries yellow halo
Candida tropicalis 750	50-100	luxuriant	>=50%	smooth discrete, dark brown with black centres, diffused blackening after 72hrs, sheen, slight mycelial fringe
Streptococcus pyogenes 19615	50-100	luxuriant	>=50%	-
Escherichia coli 25922	>=10 ³	inhibited	0%	-
Staphylococcus aureus 25923	>=10 ³	inhibited	0%	-
Candida pseudotropicalis	50-100	Good	40-50%	Dark reddish brown, glistening colony

TECHNICAL SHEET

Precautions :	1. In Vitro diagnostic use only. 2. Read the label before opening the container
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.
Use:	For detection, selective isolation, differentiation and presumptive identification of <i>Candida albicans</i> and <i>Candida Tropicalis</i> .
Storage:	Store between 15-25°C. Use before expiry date on the label.
Packing:	20/50 disposable plates.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.