

TECHNICAL SHEET

PP030	TCBS Agar Plate				
Formula					
Ingredients:		gms/lit.			
Proteose peptone		10.00			
Yeast extract		5.00			
Sodium thiosulphate		10.00			
Sodium citrate		10.00			
Bile		8.00			
Sucrose		20.00			
Sodium chloride		10.00			
Ferric citrate		1.00			
Bromo thymol blue		0.04			
Thymol blue		0.04			
Agar		15.00			
Final pH (at 25°C): 8.6 ± 0.2					
Directions:					
Label the ready to use plate (PP030). Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.					
Principle:					
Proteose peptone and yeast extract provide nitrogenous compounds, vitamin B complex and other essential growth nutrients. Bile, a derivative of bile salts and sodium citrate inhibit gram-positive bacteria and coliforms. Sodium thiosulphate serves as a good source of sulphur, which in combination with ferric citrate detects the production of hydrogen sulphide. Bromothymol blue and thymol blue are the pH indicators. For the metabolism of Vibrios, sucrose is added as a fermentable carbohydrate. Vibrio that is able to utilize sucrose will form yellow colonies. A few strains of <i>V. cholerae</i> may appear green or colorless on TCBS Agar due to delayed sucrose fermentation.					
(I) QC Tests					
	pH:	8.6 ± 0.2			
	Color:	Bluish green coloured medium.			
	Appearance:	Sterile TCBS Agar in 85mm disposable plates.			
(II) Sterility test		Passes release criteria			
(III) Q.C. Test Microbiological					
Cultural characteristics observed after incubation at 35-37°C for 18-48 hours.					
	MICROORGANISM (ATCC)	INOCULUM	GROWTH	RECOVERY	COLOR OF COLONY
	<i>Escherichia coli</i> 25922	>=10 ³	inhibited	0%	bluish green
	<i>Vibrio parahaemolyticus</i> 17802 (00037*)	50-100	good-luxuriant	>=50%	cream pink
	<i>Enterococcus faecalis</i> 29212	>=10 ³	inhibited	0%	-
	<i>Vibrio vulnificus</i> 29306	50-100	Fair-good	>=50%	greenish yellow
	<i>Vibrio fluvialis</i> 33809	50-100	Good-luxuriant	>=50%	yellow
	<i>Vibrio cholerae</i> 15748	50-100	Good-luxuriant	>=50%	yellow
	<i>Shigella flexneri</i> 12022	>=10 ³	inhibited	0%	-

Refer disclaimer Overleaf

TECHNICAL SHEET

Precautions :	1. In Vitro diagnostic use only.
	2. Read the label before opening the container
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. TCBS Agar is highly selective for Vibrio species. Any H ₂ S negative colony of TCBS Agar can be considered presumptive positive for Vibrio.
Use:	Recommended for the selective isolation of Vibrio cholerae and other enteropathogenic Vibrio's.
Storage:	Store between 20-30°C. Use before expiry date on the label.
Packing:	20/50 disposable plates.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.