BIOMARK Laboratories-INDIA www.biomarklabs.com

TECHNICAL SHEET

PP043 Potato Dextrose A	gar Plate
Formula	
Ingredients:	gms/lit.
Potatoes infusion from	200.00
Dextrose	20.00
Agar	15.00
Final pH (at 25°C): 5.6 <u>+</u> 0.2	
Directions:	
). Either streak, inoculate or surface spread the
test inoculum (50-100 CFU) aseptica	ally on the plate.
Principle:	
acidification should be avoided as i	the bacterial growth. Heating the medium after t may hydrolyze the agar which can render the
sporulation, for maintaining stock	Dextrose Agar is also used for stimulating cultures of certain dermatophytes and for of dermatophytes on the basis of pigment
sporulation, for maintaining stock differentiation of typical varieties production	c cultures of certain dermatophytes and for
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests	c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests pH:	c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment 5.6 ± 0.2
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests pH: Color:	 c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment 5.6 ± 0.2 Light Amber coloured medium Sterile Potato Dextrose Agar in 150mm disposable
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests pH: Color: Appearance:	 c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment 5.6 ± 0.2 Light Amber coloured medium Sterile Potato Dextrose Agar in 150mm disposable plates.
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests pH: Color: Appearance: (II)Sterility test (III)Q.C. Test Microbiological	 c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment 5.6 ± 0.2 Light Amber coloured medium Sterile Potato Dextrose Agar in 150mm disposable plates.
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests pH: Color: Appearance: (II)Sterility test (III)Q.C. Test Microbiological	 c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment 5.6 ± 0.2 Light Amber coloured medium Sterile Potato Dextrose Agar in 150mm disposable plates. Passes release criteria
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests pH: Color: Appearance: (II)Sterility test (III)Q.C. Test Microbiological Cultural characteristics observed	 c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment 5.6 ± 0.2 Light Amber coloured medium Sterile Potato Dextrose Agar in 150mm disposable plates. Passes release criteria after an incubation at 22 - 28°C for 48-72 hours.
sporulation, for maintaining stock differentiation of typical varieties production (I) QC Tests pH: Color: Appearance: (III)Sterility test (III)Q.C. Test Microbiological Cultural characteristics observed MICROORGANISM (ATCC)	 c cultures of certain dermatophytes and for of dermatophytes on the basis of pigment 5.6 ± 0.2 Light Amber coloured medium Sterile Potato Dextrose Agar in 150mm disposable plates. Passes release criteria after an incubation at 22 - 28°C for 48-72 hours. GROWTH luxuriant luxuriant

Refer disclaimer Overleaf

Page 01 of 02

BIOMARK Laboratories-INDIA www.biomarklabs.com TECHNICAL SHEET

Precautions :	1. In Vitro diagnostic use only.
	2. Read the label before opening the container
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.
Use:	For isolation and enumeration of yeasts and moulds from dairy and other products.
Storage:	Store between 2-8°C. Use before expiry date on the label.
Packing:	20/50 disposable plates.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

Page 02 of 02